

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 09:31:22 ON 08 NOV 2007

=> index bioscience

FILE 'DRUGMONOG' ACCESS NOT AUTHORIZED

COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
0.63	0.63

FULL ESTIMATED COST

INDEX 'ADISCTI, ADISINSIGHT, ADISNEWS, AGRICOLA, ANABSTR, ANTE, AQUALINE, AQUASCI, BIOENG, BIOSIS, BIOTECHABS, BIOTECHDS, BIOTECHNO, CABA, CAPLUS, CEABA-VTB, CIN, CONFSCI, CROPB, CROPU, DDFB, DDFU, DGENE, DISSABS, DRUGB, DRUGMONOG2, DRUGU, EMBAL, EMBASE, ...' ENTERED AT 09:33:13 ON 08 NOV 2007

69 FILES IN THE FILE LIST IN STNINDEX

Enter SET DETAIL ON to see search term postings or to view  
search error messages that display as 0\* with SET DETAIL OFF.

=> s spongilla lacustris

7	FILE AGRICOLA
2	FILE AQUALINE
86	FILE AQUASCI
1	FILE BIOENG
215	FILE BIOSIS
8	FILE BIOTECHNO
2	FILE CABA
53	FILE CAPLUS
2	FILE CONFSCI
1	FILE CROPB
1	FILE DDFB
5	FILE DGENE
2	FILE DISSABS
1	FILE DRUGB
36	FILE DRUGMONOG2
13	FILE EMBASE

29 FILES SEARCHED...

23	FILE ESBIODBASE
67	FILE GENBANK
1	FILE IFIPAT
2	FILE IMSPRODUCT
25	FILE LIFESCI
15	FILE MEDLINE
1	FILE NTIS
3	FILE OCEAN
59	FILE PASCAL
78	FILE SCISEARCH
20	FILE TOXCENTER
5	FILE USGENE
7	FILE USPATFULL
9	FILE WATER
1	FILE WPIDS
1	FILE WPINDEX

32 FILES HAVE ONE OR MORE ANSWERS, 69 FILES SEARCHED IN STNINDEX

L1 QUE SPONGILLA LACUSTRIS

=> s l1 and (dry or dried or lyophiliz#### or powder)

7 FILE AQUASCI

6 FILE BIOSIS

1 FILE CAPLUS

22 FILES SEARCHED...

4 FILE DRUGMONOG2

1 FILE IFIPAT

1 FILE LIFESCI

46 FILES SEARCHED...

2 FILE SCISEARCH

7 FILE USPATFULL

64 FILES SEARCHED...

1 FILE WPIDS

1 FILE WPINDEX

10 FILES HAVE ONE OR MORE ANSWERS, 69 FILES SEARCHED IN STNINDEX

L2 QUE L1 AND (DRY OR DRIED OR LYOPHILIZ#### OR POWDER)

=> file hits

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

11.97

12.60

FILE 'AQUASCI' ENTERED AT 09:44:32 ON 08 NOV 2007

COPYRIGHT 2007 FAO (On behalf of the ASFA Advisory Board). All rights reserved.

FILE 'USPATFULL' ENTERED AT 09:44:32 ON 08 NOV 2007

CA INDEXING COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'BIOSIS' ENTERED AT 09:44:32 ON 08 NOV 2007

Copyright (c) 2007 The Thomson Corporation

FILE 'DRUGMONOG2' ENTERED AT 09:44:32 ON 08 NOV 2007

COPYRIGHT (C) 2007 IMSWORLD Publications Ltd

FILE 'SCISEARCH' ENTERED AT 09:44:32 ON 08 NOV 2007

Copyright (c) 2007 The Thomson Corporation

FILE 'CAPLUS' ENTERED AT 09:44:32 ON 08 NOV 2007

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'IFIPAT' ENTERED AT 09:44:32 ON 08 NOV 2007

COPYRIGHT (C) 2007 IFI CLAIMS(R) Patent Services (IFI)

FILE 'LIFESCI' ENTERED AT 09:44:32 ON 08 NOV 2007

COPYRIGHT (C) 2007 Cambridge Scientific Abstracts (CSA)

FILE 'WPIDS' ENTERED AT 09:44:32 ON 08 NOV 2007

COPYRIGHT (C) 2007 THE THOMSON CORPORATION

FILE 'WPINDEX' ACCESS NOT AUTHORIZED

=> s l2

L3 30 L2

=> d rank

F1 7 AQUASCI

	ENTRY	SESSION
FULL ESTIMATED COST	62.70	75.30

FILE 'STNGUIDE' ENTERED AT 09:46:00 ON 08 NOV 2007  
USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT  
COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

FILE CONTAINS CURRENT INFORMATION.  
LAST RELOADED: Nov 2, 2007 (20071102/UP).

	SINCE FILE	TOTAL
	ENTRY	SESSION
=> log y		
COST IN U.S. DOLLARS		
FULL ESTIMATED COST	0.30	75.60

STN INTERNATIONAL LOGOFF AT 09:48:42 ON 08 NOV 2007

F2	7	USPATFULL
F3	6	BIOSIS
F4	4	DRUGMONOG2
F5	2	SCISEARCH
F6	1	CAPLUS
F7	1	IFIPAT
F8	1	LIFESCI
F9	1	WPIDS
F10	1	WPINDEX

=> dup rem l3

DUPLICATE IS NOT AVAILABLE IN 'DRUGMONOG2'.

ANSWERS FROM THESE FILES WILL BE CONSIDERED UNIQUE  
PROCESSING COMPLETED FOR L3

L4 22 DUP REM L3 (8 DUPLICATES REMOVED)  
ANSWERS '1-6' FROM FILE AQUASCI  
ANSWERS '7-13' FROM FILE USPATFULL  
ANSWERS '14-16' FROM FILE BIOSIS  
ANSWERS '17-20' FROM FILE DRUGMONOG2  
ANSWERS '21-22' FROM FILE SCISEARCH

=> d bib abs 1-6 14-16 21-22

L4 ANSWER 1 OF 22 AQUASCI COPYRIGHT 2007 FAO (On behalf of the ASFA  
Advisory Board). All rights reserved. on STN DUPLICATE 2  
AN 2002:47073 AQUASCI Full-text  
DN ASFA1 2002  
TI Life cycles and ecological interactions of freshwater sponges (Porifera,  
Spongillidae) in the river Rhine in Germany.  
AU Gugel, J.  
CS Ringstrasse 1, D-91091 Grossenseebach FRG); E-mail: j\_gugel@hotmail.co  
SO Limnologica, (20010000) vol. 31, no. 3, pp. 185-198.  
ISSN: 0075-9511.  
DT Journal  
FS ASFA1  
LA English  
SL English  
AB In the years 1993-1995 the species composition and ecological interactions of  
freshwater sponges (Porifera, Spongillidae) were investigated in the river  
Rhine between Karlsruhe and Bonn (Germany). The species *Trochospongilla*  
*horrida*, *Ephydatia fluviatilis*, *Spongilla lacustris*, *Ephydatia muelleri*,  
*Eunapius fragilis* and *Eunapius carteri* were found. *E. muelleri*, *T. horrida*, *S.*  
*lacustris* and *Eun. fragilis* were only found in habitats which remained flooded  
year-round, whereas *E. fluviatilis* occurred in similar habitats, but its  
abundance remained high at places which dried out regularly. Colonies of  
*E. fluviatilis* were found throughout the year, colonies of *E. muelleri*, *T.*  
*horrida*, *S. lacustris* and *Eun. fragilis* were only present during the warm  
season, in the cold season only gemmules of the latter species were found. The  
dispersal of *E. fluviatilis* within the habitat was mainly ensured by sexually  
produced larvae; the role of asexually produced gemmules was negligible. *Eun.*  
*fragilis* was also dispersed via larvae, it ensured its overwintering through  
production of abundant gemmules in *E. muelleri*, *T. horrida* and *S. lacustris*  
sexually produced larvae were not observed. Beside having no free moving  
larvae, the gemmules of *T. horrida* were tightly fixed to the substrate, so the  
distribution ability of this species was restricted. In the case of *E.*  
*muelleri* and *S. lacustris* some gemmules became free and were dispersed within  
the habitat. Other gemmules rested in the dead body of their mothersponges and  
ensured the recolonization of the same place in following suitable seasons.  
*Ephydatia muelleri* produced many gemmules, using almost its whole tissue for

gemmulation before desintegrating. Freshwater sponges in general interact with a number of animals from other phyla, nevertheless these associations were less frequent as reported in the literature at other places.

ANSWER 2 OF 22 AQUASCI COPYRIGHT 2007 FAO (On behalf of the ASFA Advisory Board). All rights reserved. on STN DUPLICATE 3

AN 82:3786 AQUASCI Full-text

DN ASFA1 1982 12-14169

TI Population Dynamics and Standing Biomass of the Freshwater Sponge *Spongilla lacustris* .

AU Frost, T.M.; DeNagy, G.S.; Gilbert, J.J.

CS Lab. Limnol., Univ. Wisconsin, Madison, WI 53706, USA

SO ECOLOGY., (1982) vol. 63, no. 5, pp. 1203-1210.

DT Journal

FS ASFA1

LA English

SL English

AB In a sphagnum bog-pond in central New Hampshire, USA, the common freshwater sponge *S. lacustris* exhibits a high growth rate and reaches a large standing biomass (ash-free dry biomass is 1.7 g/m super(2)) by the end of its growing season in October. The sponge overwinters as gemmules, and the conversions between these dormant forms and active sponge tissue in the spring and fall are very conservative in terms of biomass. *S. lacustris* does not require a hard substratum. The greatest part of this population grows as upright branches from the pond bottom or attached to aquatic macrophytes. In winter, the gemmulated sponges, including those attached to macrophytes, collapse into the bottom sediments. There is a high mortality during this period, resulting from the accumulation of sediments over the gemmulated sponges, which prevents their successful hatching. It appears that sponges derived from sexually produced larvae would be particularly unlikely to survive the winter. Asexual growth processes are thus of primary quantitative importance in this system and must be considered in evaluating reproductive effort.

L4 ANSWER 3 OF 22 AQUASCI COPYRIGHT 2007 FAO (On behalf of the ASFA Advisory Board). All rights reserved. on STN DUPLICATE 4

AN 80:12843 AQUASCI Full-text

DN ASFA1 1981 11-09065

TI Clearance rate determinations for the freshwater sponge *Spongilla lacustris* : effects of temperature, particle type and concentration, and sponge size.

AU Frost, T.M.

CS Dep. EPO Biol., Univ. Colorado, Boulder, CO 80309, USA

SO Arch. Hydrobiol., (1980) Vol. 90, No. 3, pp. 330-356.

DT Journal

FS ASFA1

LA English

SL English

AB The clearance rates for *S. lacustris* on a single particle type, the bacterium *Aerobacter aerogenes* or simultaneously on *A. aerogenes* and the yeast *Rhodotorula glutinis* or the alga *Chlamydomonas reinhardtii* were determined. For most particle densities, there is no relationship between cell concentration and clearance rate. There is an effect of sponge size and water temperature on clearance rates and these relationships are described using quadratic regression equations. In general, there is a good correlation between clearance rates determined simultaneously on two particle types. However, when *R. glutinis* is present at a high density, clearance rates on it may be drastically lowered while they are maintained at a nearly normal level for *A. aerogenes* present at lower concentrations. It appears that selection of

particles can occur in the digestive activities of the sponge. This selection, facilitated by the parazoan nature of the sponge digestive system, is accomplished by a means unavailable to other groups of suspension feeders. The clearance rates normally exhibited by *S. lacustris* are comparable to those of zooplankton on a per dry weight and on a total community basis.

L4 ANSWER 4 OF 22 AQUASCI COPYRIGHT 2007 FAO (On behalf of the ASFA  
Advisory Board). All rights reserved. on STN DUPLICATE 5  
AN 2002:10906 AQUASCI Full-text  
DN ASFA1 2002  
TI Chlorophyll and primary productivity of some green, freshwater sponges.  
AU Gilbert, J.J.; Allen HL  
CS Dep. Biol. Sci. Dartmouth Coll., Hanover, NH, USA  
SO Int. Rev. Gesamten Hydrobiol., (19730000) vol. 58, no. 5, pp. 633-638..  
Records keyed from 1975 ASFA printed journals..  
DT Journal  
FS ASFA1  
LA English  
AB Relationships between ash-free dry wt, viable chlorophyll a and pheopigments a  
were examined by linear regression analysis for non-gemmulating and  
gemmulated *Spongilla lacustris* and for an unidentified, non-gemmulating sp.  
The primary productivity of *S. lacustris* was determined by both O Sub(2) and  
Super(14)C techniques and expressed on a biomass, chlorophylla. and area  
basis, and the photosynthetic efficiency was calculated. The proportion of  
algal photosynthate excreted by the sponge was investigated, and the transfer  
of algal photosynthate to the sponge was examined by autoradiography.

L4 ANSWER 5 OF 22 AQUASCI COPYRIGHT 2007 FAO (On behalf of the ASFA  
Advisory Board). All rights reserved. on STN  
AN 84:22450 AQUASCI Full-text  
DN ASFA1 1986 16-05216  
TI Macrofauna inhabiting the colonies of the sponge *Spongilla*  
*lacustris* (L.) in the River Gac.  
CONGRESS IN FRANCE 1983. PROCEEDINGS.  
AU Konopacka, A.; Sicinski, J.  
CS Univ. Lodz, Dep. Gen. Zool., Lodz, Poland  
SO VERH. INT. VER. THEOR. ANGEW. LIMNOL./PROC. INT. ASSOC. THEOR. APPL.  
LIMNOL./TRAV. ASSOC. INT. LIMNOL. THEOR. APPL., (1984) pp. 2968-2973.  
Meeting Info.: (22.) Congress of the International Association of  
Limnology. (Lyon (France)). 21 Aug 1983.  
ISSN: 0368-0770 ISBN: 3-510-54025-5.  
DT Book  
TC Conference  
FS ASFA1  
LA English  
SL English  
AB Colonies of the freshwater sponge, *Spongilla lacustris*, were chosen for  
studies of the fauna inhabiting sponge tissues. Samples of these tissues were  
regularly taken from February to November 1979; the numbers of animals were  
~~calculated per 10g of the dry sponge tissue weight.~~ Among the groups found,  
*Chironomidae* and *Oligochaeta* clearly dominated. Their summarized percentage  
share in all months was usually higher than 70% of all individuals inhabiting  
sponges. Changes in the abundance of the sponge in fauna in the yearly cycle  
were caused primarily by the phenology of particular species and to a lesser  
extent by the presence or absence of the liver sponge tissue.

L4 ANSWER 6 OF 22 AQUASCI COPYRIGHT 2007 FAO (On behalf of the ASFA

Advisory Board). All rights reserved. on STN

AN 78:6856 AQUASCI Full-text  
DN ASFA1 1979 09-02117  
TI The relationship of light duration to oxygen consumption in the green, freshwater sponge *Spongilla lacustris*.  
AU Szuch, E.J.; Studier, E.H.; Sullivan, R.B., Jr.  
CS Dep. Biol., Univ. Michigan-Flint, Flint, MI 48503, USA  
SO Comp. Biochem. Physiol., (1978) Vol. 50A, No. 2, pp. 221-223.  
DT Journal  
FS ASFA1  
LA English  
SL English  
AB Oxygen consumption of green *Spongilla lacustris* in the light is lower than that in the dark. The relationship between oxygen consumption (OC in  $\mu\text{L O}_2/\text{mg dry wt/hr}$ ) and hours of continuous light (HL) is  $\text{OC} = 1.207 - 0.0143 (\text{HL})$  SUP-2. For light durations greater than 9 hr, symbiotic zoochlorellae produce photosynthetically derived oxygen in excess of the needs of the symbiotic complex.

L4 ANSWER 14 OF 22 BIOSIS COPYRIGHT (c) 2007 The Thomson Corporation on STN

AN 1987:103717 BIOSIS Full-text  
DN PREV198783052695; BA83:52695  
TI OXYGEN CONSUMPTION OF FRESHWATER SPONGES.  
AU KHARCHENKO T A [Reprint author]; LYASHENKO A V  
CS INST HYDROBIOL, ACAD SCI UKR SSR, KIEV, USSR  
SO *Gidrobiologicheskii Zhurnal*, (1986) Vol. 22, No. 3, pp. 98-100.  
CODEN: GBZUAM. ISSN: 0375-8990.

DT Article  
FS BA  
LA RUSSIAN  
ED Entered STN: 26 Feb 1987  
Last Updated on STN: 26 Feb 1987

AB Oxygen consumption by fresh-water *Spongilla lacustris* L. is established to depend on the magnitude of dry weight of colonies. The amount of organic matters mineralized by colonies of this sponge in benthos and periphyton of the main part of the Dnieper-Donbas [USSR] channel is calculated.

L4 ANSWER 15 OF 22 BIOSIS COPYRIGHT (c) 2007 The Thomson Corporation on STN

AN 1978:258825 BIOSIS Full-text  
DN PREV197866071322; BA66:71322  
TI THE RELATIONSHIP OF LIGHT DURATION TO OXYGEN CONSUMPTION IN THE GREEN FRESH WATER SPONGE *SPONGILLA-LACUSTRIS*.  
AU SZUCH E J [Reprint author]; STUDIER E H; SULLIVAN R B JR  
CS DEP BIOL, UNIV MICH, FLINT, MICH 48503, USA  
SO Comparative Biochemistry and Physiology A, (1978) Vol. 60, No. 2, pp. 221-224.  
CODEN: CBPAB5. ISSN: 0300-9629.

DT Article  
FS BA  
LA ENGLISH

AB  $\text{O}_2$  consumption of green *S. lacustris* in the light is lower than that in the dark. The relationship between  $\text{O}_2$  consumption (OC in  $\mu\text{L O}_2/\text{mg dry wt per h}$ ) and hours of continuous light (HL) is  $\text{OC} = 1.207 - 0.0143 (\text{HL})^2$ . For light durations  $> 9$  h, symbiotic zoochlorellae produce photosynthetically derived  $\text{O}_2$  in excess of the needs of the symbiotic complex.

L4 ANSWER 16 OF 22 BIOSIS COPYRIGHT (c) 2007 The Thomson Corporation on  
STN

AN 1927:11492 BIOSIS Full-text

DN PREV19270100013749; BA01:13749

TI Structure and life of German freshwater sponges with a key to forms  
hitherto described from Europe.  
Original Title: Bau und Leben der deutschen Siisswasserschwamme. Mit einer  
Bestimmungstabelle der bisher in Europa gefundenen Formen.

AU ARNDT, WALTHER

SO MIKROSK F NATURFREUNDE, (1926) Vol. 4, No. 5, pp. 119-132.

DT Article

FS BA

LA Unavailable

ED Entered STN: May 2007  
Last Updated on STN: May 2007

AB Introduction to the anatomy and physiology of the Spongillinae of Germany,  
including an enumeration of the fossil species. Habitus-photographs of  
certain forms are given. Spongilla lacustris has oscular "chimneys"  
projecting 3/4-1 cm. which contract to a barely perceptible hump when  
stimulated; the reaction time for this movement was found to be 5-10 min.  
Chemical analysis of dried S. lacustris showed a relatively high iron content.  
Facts are given concerning the use of fresh-water sponges in medicine. The  
technique employed in determining spongillid species is explained. The key  
covers the known forms (28) of Europe, including fossil species. ABSTRACT  
AUTHORS: W. Arndt (transl. by M. W. de Laubenfels)

  

L4 ANSWER 21 OF 22 SCISEARCH COPYRIGHT (c) 2007 The Thomson Corporation on  
STN

AN 1997:632480 SCISEARCH Full-text

GA The Genuine Article (R) Number: XR976

TI Particle capture and deposition by deep sea sponges from the  
Norwegian-Greenland Sea

AU Witte U (Reprint); Brattegard T; Graf G; Springer B

CS GEOMAR RES CTR, DEPT ENVIRONM GEOL, WISCHHOFSTR 1-3, D-24148 KIEL, GERMANY  
(Reprint); UNIV BERGEN, DEPT FISHERIES & MARINE BIOL, N-5020 BERGEN,  
NORWAY; UNIV ROSTOCK, DEPT MARINE BIOL, D-18055 ROSTOCK, GERMANY

CYA GERMANY; NORWAY

SO MARINE ECOLOGY-PROGRESS SERIES, (1997) Vol. 154, pp. 241-252.  
ISSN: 0171-8630.

PB INTER-RESEARCH, NORDBUNTE 23, D-21385 OLDENDORF LUHE, GERMANY.

DT Article; Journal

FS AGRI

LA English

REC Reference Count: 57

ED Entered STN: 1997  
Last Updated on STN: 1997  
\*ABSTRACT IS AVAILABLE IN THE ALL AND IALL FORMATS\*

AB Particle uptake and deposition by the 2 most abundant deep-sea  
demosponge species from the Norwegian and Greenland Sea (Thenea  
abyssorum) and the deep fjords of western Norway (Thenea muricata) were  
studied in flume experiments. Fluorescent particles of 1, 2, 3, 6, 10  
and 16  $\mu\text{m}$  diameter (microspheres, Duke Scientific Corporation(R)) with  
a density of  $1.05\text{ g cm}^{-3}$  were used at 2 current velocities, 1.5 and 5  
 $\text{cm s}^{-1}$ . Both species ingested small particles exclusively ( $<6\text{ }\mu\text{m}$   
and  $<10\text{ }\mu\text{m}$ , respectively), with a preference for the smallest fraction  
at both current speeds. The results suggest that the size spectrum  
actually ingested depends on the supplied particle sizes rather than on  
current velocity. Current velocity fields around dead specimens were



recorded and turbulence intensity calculated in order to determine the influence of the sponge acting as a biogenic structure on the near-bottom current regime. Disruption of flow conditions was detected as far as 14 cm downstream and several cm laterally from the biogenic obstacles. Bulk biodeposition rates calculated from sponge biomass and volume of ingested particles range between 7 and 10 mg d<sup>(-1)</sup> g<sup>(-1)</sup> ash-free dry weight.

L4 ANSWER 22 OF 22 SCISEARCH COPYRIGHT (c) 2007 The Thomson Corporation on STN

AN 1993:425863 SCISEARCH Full-text

GA The Genuine Article (R) Number: LL063

TI SUSPENSION-FEEDING IN MARINE SPONGES HALICHONDRIA-PANICEA AND HALICLONA-URCEOLUS - EFFECTS OF TEMPERATURE ON FILTRATION-RATE AND ENERGY-COST OF PUMPING

AU RIISGARD H U.(Reprint); THOMASSEN S; JAKOBSEN H; WEEKS J M; LARSEN P S

CS ODENSE UNIV, INST BIOL, DK-5230 ODENSE, DENMARK (Reprint); TECH UNIV DENMARK, DEPT FLUID MECH, DK-2800 LYNGBY, DENMARK

CYA DENMARK

SO MARINE ECOLOGY-PROGRESS SERIES, (JUN 1993) Vol. 96, No. 2, pp. 177-188. ISSN: 0171-8630.

PB INTER-RESEARCH, NORDBUNTE 23, D-21385 OLDENDORF LUHE, GERMANY.

DT Article; Journal

FS AGRI

LA English

REC Reference Count: 37

ED Entered STN: 1994

Last Updated on STN: 1994

\*ABSTRACT IS AVAILABLE IN THE ALL AND IALL FORMATS\*

AB Filtration rate (measured as clearance of algal cells) was measured at different temperatures in the sponge Halichondria panicea. An increase in water temperature from 6 to 12-degrees-C caused the mean filtration rate to increase 4.3 +/- 2.3 times. This value was higher than previously found for other marine ciliary suspension-feeding animals. Filtration rate at 12-degrees-C was also measured in Haliclona urceolus by means of an indirect clearance method in addition to a direct technique for measuring pumping rate. It was found that the 2 sponge species had near-identical filtration rates, with maximum rates of approximately 60 ml min<sup>-1</sup> (g dry weight)<sup>-1</sup> at 12-degrees-C. The normal pump pressure, or operating point O(p), of a standard sponge (based on our own measurements and calculations from literature data for a 0.1 g dry weight Haliclona sp.) was estimated as the sum of main contributions to head losses along the flow path from entry (ostia) to exit (osculum). The head losses were as follows: ostia 0.0373 mm H<sub>2</sub>O; inhalant canal 0.1205 to 0.013 mm H<sub>2</sub>O; prosopyles 0.1153 to 0.02321 mm H<sub>2</sub>O; collar-filter 0.122 mm H<sub>2</sub>O; exhalant canals = inhalant canals; and osculum 0.1576 mm H<sub>2</sub>O. The (maximal) O(p) was found to be 0.673 mm H<sub>2</sub>O and the power output P(p) from the sponge pump was 0.677 muW. The pump work, defined as P(p)R<sup>-1</sup> where R is the respiratory output, was 0.85 %. The low energy cost of filtration and the temperature effect are discussed and compared with recent data for other ciliary suspension feeders. It is argued that passive current-induced filtration may be of insignificant importance for sponges.

=> file stnguide

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

# ● WEST Search History ●

DATE: Thursday, November 08, 2007

Hide?	<u>Set Name</u>	<u>Query</u>	<u>Hit Count</u>
		<i>DB=PGPB,USPT; PLUR=YES; OP=ADJ</i>	
<input type="checkbox"/>	L3	11 same (powder dry dried lyophiliz\$4)	1
		<i>DB=EPAB,JPAB,DWPI; PLUR=YES; OP=ADJ</i>	
<input type="checkbox"/>	L2	spongilla lacustris	0
		<i>DB=PGPB,USPT; PLUR=YES; OP=ADJ</i>	
<input type="checkbox"/>	L1	spongilla lacustris	9

END OF SEARCH HISTORY

## Case Creation Option

*Case "10659451us20071108" already exists. Please overwrite it or cancel the operation.*

### The Contents of Case "10659451us20071108"

Qnum	Query	DB Name	Thesaurus	Operator	Plural
Q1	7241871.pn.	USPT	None	ADJ	YES
Q2	fresh watert or freshwater	USPT	None	ADJ	YES
Q3	sponge	USPT	None	ADJ	YES
Q4	spongilla lacustris	USPT	None	ADJ	YES
Q5	powder or solids	USPT	None	ADJ	YES
Q6	Q2 near5 Q3	USPT	None	ADJ	YES
Q7	fresh water or freshwater	USPT	None	ADJ	YES
Q8	Q3 near5 Q7	USPT	None	ADJ	YES
Q9	Q5 near5 Q8	USPT	None	ADJ	YES
Q10	Q5 and Q8	USPT	None	ADJ	YES
Q11	Q4 near5 Q3	USPT	None	ADJ	YES
Q12	Q5 near5 Q11	USPT	None	ADJ	YES
Q13	Q10 and Q7	USPT	None	ADJ	YES
Q14	Q3 and Q13	USPT	None	ADJ	YES
Q15	Q4 and Q14	USPT	None	ADJ	YES
Q16	collagen or protein	USPT	None	ADJ	YES
Q17	dry or lyophilized or (freeze-dried)	USPT	None	ADJ	YES
Q18	Q16 same Q17	USPT	None	ADJ	YES
Q19	Q4 same Q7	USPT	None	ADJ	YES
Q20	Q3 same Q7	USPT	None	ADJ	YES
Q21	Q18 same Q20	USPT	None	ADJ	YES
Q22	Q18 and Q20	USPT	None	ADJ	YES

Q23	Q2 and Q20	USPT	None	ADJ	YES
Q24	Q4 and Q20	USPT	None	ADJ	YES
Q25	Q16 and Q20	USPT	None	ADJ	YES
Q26	coollagen	USPT	None	ADJ	YES
Q27	collagen	USPT	None	ADJ	YES
Q28	Q20 and Q27	USPT	None	ADJ	YES
Q29	Q4 and Q27	USPT	None	ADJ	YES
Q30	Q4 with Q27	USPT	None	ADJ	YES
Q31	Q4 same Q27	USPT	None	ADJ	YES
Q32	Q4 near5 Q27	USPT	None	ADJ	YES
Q33	Q4 near5 Q17	USPT	None	ADJ	YES
Q34	Q4 and Q17	USPT	None	ADJ	YES
Q35	Q4 same Q17	USPT	None	ADJ	YES
Q36	(jojoba oil)	USPT	None	ADJ	YES
Q37	Q34 and Q36	USPT	None	ADJ	YES
Q38	Q4 and Q36	USPT	None	ADJ	YES
Q39	Q3 and Q36	USPT	None	ADJ	YES
Q40	Q8 and Q36	USPT	None	ADJ	YES
Q41	("flora Ester 60") or (flora ester) or(hohoba oil) or lipovol or (jojoba oil) or ("RN-61789-91-1")	USPT	None	ADJ	YES
Q42	Q4 same Q17	USPT	None	ADJ	YES
Q43	dried or (freeze dried) or powdered	USPT	None	ADJ	YES
Q44	Q4 same Q43	USPT	None	ADJ	YES
Q45	Q4 near5 Q43	USPT	None	ADJ	YES
Q46	Q4 near10 Q43	USPT	None	ADJ	YES
Q47	Q4 and Q43	USPT	None	ADJ	YES
Q48	Q41 and Q47	USPT	None	ADJ	YES
Q49	Q4 and Q41	USPT	None	ADJ	YES
Q50	insoluble components	USPT	None	ADJ	YES

Q51	asntiseptic or gel or (enzyme gel) or emolient or fragrance or (boric acid)	USPT	None	ADJ	YES
Q52	Q50 and Q4	USPT	None	ADJ	YES
Q53	Q51 and Q4	USPT	None	ADJ	YES
Q54	Q47 and Q53	USPT	None	ADJ	YES
Q55	sponge and (freshwater) or (Fresh water)	PGPB,EPAB,JPAB,DWPI	None	ADJ	YES
Q56	(spongilla lacustris)	PGPB,EPAB,JPAB,DWPI	None	ADJ	YES
Q57	dried or (freeze dried) or powdered or dry or lyophilized	PGPB,EPAB,JPAB,DWPI	None	ADJ	YES
Q58	("flora Ester 60") or (flora ester) or(hohoba oil) or lipovol or (jojoba oil) or ("RN-61789-91-1")	PGPB,EPAB,JPAB,DWPI	None	ADJ	YES
Q59	asntiseptic or gel or (enzyme gel) or emolient or fragrance or (boric acid)	PGPB,EPAB,JPAB,DWPI	None	ADJ	YES
Q60	Q55 and Q56	PGPB,EPAB,JPAB,DWPI	None	ADJ	YES
Q61	Q57 and Q56	PGPB,EPAB,JPAB,DWPI	None	ADJ	YES
Q62	Q58 and Q61	PGPB,EPAB,JPAB,DWPI	None	ADJ	YES
Q63	Q59 and Q61	PGPB,EPAB,JPAB,DWPI	None	ADJ	YES
Q64	20030211989.pn.	PGPB	None	ADJ	YES
Q65	Q64 and @pd > 20071106	PGPB	None	ADJ	YES
Q66	Spongilla lacustris	PGPB,USPT	None	ADJ	YES
Q67	Spongilla lacustris	EPAB,JPAB,DWPI	None	ADJ	YES
Q68	Q66 same (powder dry dried lyophiliz\$4)	PGPB,USPT	None	ADJ	YES

---

 Overwrite

Cancel